



Sensitivity of tropical water and energy cycle to SST increase and doubling CO₂ as simulated with an upgraded Multiscale Modeling Framework



Kuan-Man Xu¹ , Anning Cheng²

1. NASA Langley Research Center, Hampton, VA
2. NOAA/EMC, College Park, MD



Outline

1. Why focus on low clouds?
2. Model description and simulation setup
3. Results from control and sensitivity simulations
4. Comparison with earlier studies
5. Summary and conclusions



Objectives



- ✦ to improve the simulation of low-level clouds using a multiscale modeling framework (MMF) with a third-order turbulence closure in its CRM component
- ✦ to understand climate sensitivity and cloud response from this MMF
- ✦ to compare with studies using an MMF with a simple first-order turbulence closure

Cheng and Xu (2011; *JGR*); Xu and Cheng (2013a,b; *J. Climate*)

Cheng and Xu (2013a; *J. Climate*); Cheng and Xu (2013b; *JGR*)



Cheng and Xu (2014, *JGR*); Painemal et al. (2015; *J. Climate*)

Cheng and Xu (2015, *J. Climate*)




Outline

1. Why focus on low clouds?
2. Model description and simulation setup
3. Results from control and sensitivity simulations
4. Comparison with earlier studies
5. Summary and conclusions



Uncertainties in cloud feedback remain in GCMs


Local contribution to intermodel spread in cloud feedback: AR4



- Most of intermodel spread arises from low stratocumulus/cumululus regions

Soden and Vecchi (2011)

Local contribution to intermodel spread in cloud feedback: AR5





- Low subtropical clouds still uncertain.
- Large contribution from equatorial Pacific.

Soden and Vecchi (2011):

- Low cloud cover is responsible for $\sim 3/4$ of the difference in global-mean net cloud feedback among AR4 models, with the largest contributions associated with low-level subtropical marine cloud systems;



The low-cloud inconsistency and deficiency in most of the models.




SE Pacific Stratocumulus

Subtropical stratocumulus




4 September 2009 at 20:45 UTC



from Wood (2012; *Mon. Wea. Rev.*)





Processes associated with stratocumulus




Outline

1. Why focus on low clouds?
2. Model description and simulation setup
3. Results from control and sensitivity simulations
4. Comparison with earlier studies
5. Summary and conclusions




IPHOC: Intermediately-prognostic higher-order turbulence closure for cloud-resolving model (CRM)



The Multiscale Modeling Framework

(Grabowski 2001; Khairoutdinov and Randall 2001; Cheng & Xu 2011; Xu and Cheng 2013a)

- ✦ A CRM is embedded at each grid column (~100s km) of the host GCM to represent cloud physical processes
- ✦ The CRM explicitly simulates cloud-scale dynamics (~1s & km) and processes
- ✦ Periodic lateral boundary condition for CRM (not extended to the edges)





Upgraded CRM with a third-order turbulence closure (IPHOC):

- ✦ Double-Gaussian distribution of liquid-water potential temperature (θ_l), total water mixing ratio (q_t) and vertical velocity (w), pdf = $a G_1(w, q_t, \theta_l) + (1-a) G_2(w, q_t, \theta_l)$
- ✦ Skewnesses, i.e., the three third-order moments, predicted
- ✦ All first-, second-, third- and fourth-order moments, subgrid-scale condensation and buoyancy based on the same PDF
- ✦ Advantages over CLUBB (Golaz *et al.*)

1) two extra third-order moments



2) Widths of two Gaussian unequal

3) Merge to a single Gaussian if $sk = 0$





SPCAM-IPHOC climate simulations

- **SPCAM-IPHOC**
 - CAM3.5 with finite-volume dynamic core as the host GCM
 - 2-D version of System for Atmospheric Modeling (SAM) CRM with **IPHOC**
 - The CRM grid spacing is 4 km, with 32 columns, within a GCM grid box
 - The GCM grid spacing is $1.9^{\circ} \times 2.5^{\circ}$ with **32 vertical levels** (12 below 700 hPa)
- **Control simulation**
 - Forced with climatological SST and sea ice distributions (not an AMIP-type simulation) with present-day CO_2 concentration
 - Simulation duration is 10 years and 3 months, with last nine years analyzed
- **Doubled CO_2 ($2\times\text{CO}_2$) simulation**
 - Same as the control except that CO_2 is doubled; SST and sea ice are fixed
- **+2 K SST (12K) simulation**
 - Same as the control except that sea surface temperature is increased by 2 K




Outline


1. Why focus on low clouds?
2. Model description and simulation setup
3. Results from control and sensitivity simulations
4. Comparison with earlier studies
5. Summary and conclusions




Low-level (surface -- 700 hPa) cloud fraction (%) from 10-yr control simulation and changes in sensitivity simulations




More in cumulus regions in 12K; but stratocumulus regions in 2xCO₂
More in poleward of storm track in 12K, but equatorward in 2xCO₂




Longwave Cloud Radiative Forcing from 10-yr control simulation and changes in sensitivity simulations




Changes in LW CRE are larger in 12K (higher clouds due to higher SSTs); Storm tracks moves polarwards and stronger SPCZ in 12K




Surface Precipitation Rate from 10-yr control simulation and changes in sensitivity simulations




Increases at ITCZ and SPCZ (polarward movements) in I2K ;
General small decreases in tropics and over lands in $2\times\text{CO}_2$



Shortwave Cloud Radiative Forcing from 10-yr control simulation and changes in sensitivity simulations





Changes in SW CRE reflect mostly the changes in low cloud fraction in both sensitivity simulations; some due to high clouds




Outline

1. Why focus on low clouds?
2. Model description and simulation setup
3. Results from control and sensitivity simulations
4. Comparison with earlier studies
5. Summary and conclusions




Low-level (surface -- 700 hPa) cloud fraction (%) changes in sensitivity simulations (Bretherton et al. 2014)





Differences appear in cumulus regions in +SST simulations
There are large differences in CO_2 simulations everywhere

NASA



Surface Precipitation Rate changes in sensitivity simulations (Bretherton *et al.*, 2014)




Similar increases at ITCZ and SPCZ in +SST simulations;
More increases over lands in $2\times\text{CO}_2$ than in $4\times\text{CO}_2$

Changes in the tropical land and ocean (30° N/S) from CO₂ sensitivity simulations



Number in Green 2×CO₂ from
SPCAM-IPHOC multiplied by 2
Red number from 4×CO₂ in
Bretherton *et al.* (2014)

Deep convection is
enhanced over land.




Summary and conclusions

- The low cloud fractions increases in the tropics/subtropics in both sensitivity simulations, but for different reasons.
 - Enhanced inversion strength due to free tropospheric warming from 2XCO_2 ;
 - Increased instability results in more shallow convection in +2K SST
- The polarward displacement of storm tracks results in large changes (+ve polarward and -ve equatorward) in cloud fraction and cloud radiative effects in +2K SST simulation
- Precipitation increases in +2K SST are pronounced, especially ITCZ and SPCZ, but almost no increases in the global mean of 2xCO_2 simulation, in which the local changes are dis-similar to those of +2K SST in most continents
- There are many similarities with the MMF without the higher-order turbulence closure, but there are some differences
 - shallow cumulus increase more over ocean
 - stronger convection over lands and weaker convection over ocean
 - stronger lower tropospheric moisture transport over land is the reason



Temperature differences at surface and 850 hPa between sensitivity and control simulations



Increased stability in the stratocumulus regions of $2\times CO_2$
But the cumulus regions in $I2K$ do not show much change

NASA

